

BIBLIOGRAPHIE

- [1] Hérán F, Lafitte F. Imagerie en ophtalmologie pour les radiologues. In : *Œil, orbite, vision, oculomotricité*. Issy-les-Moulineaux : Elsevier Masson; 2018. p. 304.
- [2] Vignal-Clermont C, Miléa D, Tilikete C. Neuro-ophtalmologie. 2^e éd. Issy-les-Moulineaux : Elsevier Masson; 2016. p. 320.
- [3] Hérán F, Lafitte F. I.R.M. pratique en neuroradiologie. 2^e éd. Issy-les-Moulineaux : Elsevier Masson; 2013. p. 496.
- [4] Hérán F, Koskas P, Vignal C. Nerf optique. *Encycl Méd Chir. In : Ophtalmologie*. Paris : Elsevier; 2010. 21-008-A-10.
- [5] Rybicki FJ. Fast three-point Dixon MR imaging of the retrobulbar space with low-resolution images for phase correction: comparison with fast spin-echo inversion recovery imaging. *AJNR Am J Neuroradiol* 2001;22:1798–802.
- [6] Lafitte F. Imagerie de l'orbite. *Encycl Méd Chir. In : Ophtalmologie*. 10. Paris : Elsevier; 2013. p. 1–24. 21-050-A-10.
- [7] Hodel J, Rodallec M, Gerber S, et al. Séquences IRM «SWAN, SWI et VenobOLD» exploitant le phénomène de susceptibilité magnétique : principes techniques et applications cliniques. *J Neuroradiol* 2012;39:71–86.
- [8] Rott D, Leibowitz D. Optic nerve head drusen mimicking papilledema and malignant hypertension. *Eur J Intern Med* 2009;20:e112–3.
- [9] Henderson AD, Miller NR. Carotid-cavernous fistula: current concepts in aetiology, investigation, and management. *Eye (Lond)* 2018;32:164–72.
- [10] Yoshino M, Abhinav K, Yeh FC, et al. Visualization of cranial nerves using high-definition fiber tractography. *Neurosurgery* 2016;79:146–65.
- [11] Bertani GA, Bertulli L, Scola E, et al. Optic radiation diffusion tensor imaging tractography: an alternative and simple technique for the accurate detection of Meyer's loop. *World Neurosurg* 2018;117:e42–56.
- [12] Heran F, Vignal F, Lafitte F, et al. Imagerie des urgences absolues et relatives en neuro-ophtalmologie. *Encycl Méd Chir. In : Radiologie et imagerie médicale : musculosquelettique–neurologique–maxillofaciale*. Paris : Elsevier; 2016. 31-690-A-10.
- [13] Lennon VA. IgG marker of optic-spinal multiple sclerosis binds to the aquaporin-4 water channel. *J Exp Med* 2005;202:473–7.
- [14] Kidd DP, Burton BJ, Graham EM, Plant GT. Optic neuropathy associated with systemic sarcoidosis. *Neurol Neuroimmunol Neuroinflamm* 2016;3:e270.
- [15] Wingerchuk DM. The spectrum of neuromyelitis optica. *Lancet Neurol* 2007;6:805–15.
- [16] Kitley J. Longitudinally extensive transverse myelitis with and without aquaporin 4 antibodies. *JAMA Neurol* 2013;70:1375–81.
- [17] Kister I. Ultrahigh-Field MR (7 T) Imaging of brain lesions in neuromyelitis optica. In: *Mult Scler Int* 2013;2013.398259.
- [18] HO Jin K, Friedmann P, Marco ALP, et al. MRI characteristics of neuromyelitis optica spectrum disorder: an international update. *Neurology* 2015;84:1165–73.
- [19] Mukhi SV, Lincoln CM. MRI in the evaluation of acute visual syndromes. *Top Magn Reson Imaging* 2015;24:309–24.
- [20] Urolagin SB, Kotrashetti SM, Kale TP, Balihallimath LJ. Traumatic optic neuropathy after maxillofacial trauma: a review of 8 cases. *J Oral Maxillofac Surg* 2012;70:1123–30.
- [21] Kwickström P, Lindblom B, Bergström G, Zetterberg M. Amaurosis fugax: risk factors and prevalence of significant carotid stenosis. *Clin Ophthalmol* 2016;10:2165–70.
- [22] Rougier MB. Diagnosing bilateral papilledema. *J Fr Ophtalmol* 2010;33:424–9.
- [23] Lamirel C, Cassereau J, Cochereau I, et al. Papilloedema and MRI enhancement of the prechiasmal optic nerve at the acute stage of Leber hereditary optic neuropathy. *J Neurol Neurosurg Psychiatry* 2010;81:578–80.
- [24] Woodall MN, Nguyen KD, Alleyne Jr CH, Macomson SD. Bilateral transverse sinus stenosis causing intracranial hypertension. In: *BMJ Case Rep* 2013;2013.
- [25] Riggeal BD, Bruce BB, Saindane AM, et al. Clinical course of idiopathic intracranial hypertension with transverse sinus stenosis. *Neurology* 2013;80:289–95.
- [26] Manfrè L, Vero S, Focarelli-Barone C, Lagalla R. Bitemporal pseudohemianopia related to the "tilted disk" syndrome: CT, MR, and fundoscopic findings. *AJNR* 1999;20:1750–1.
- [27] Dae SL, Hee KY, Jae HK, Jeong-Min H. Morphometry of the trochlear nerve and superior oblique muscle volume in congenital superior oblique palsy. *Invest Ophthalmol Vis Sci* 2014;55:8571–5.
- [28] Grant 3rd JH, Patrinely JR, Weiss AH, et al. Trapdoor fracture of the orbit in a pediatric population. In: *Plast Reconstr Surg*. 109;2002. p. 482–9. discussion 490-5.
- [29] Saito R, Sugawara T, Mikawa S, et al. Pupil-sparing oculomotor nerve paresis as an early symptom of unruptured internal carotid-posterior communicating artery aneurysms: three case reports. *Neurol Med Chir (Tokyo)* 2008;48:304–6.
- [30] Rosenkranz M, Eckert B, Niesen WD, et al. Horner syndrome related to ipsilateral carotid wall hematoma after stent placement for the treatment of carotid stenoses. *AJNR Am J Neuroradiol* 2003;24:1508–11.
- [31] Besselmann M, Vennemann B, Lowens S, Ringelstein EB, Stogbauer F. Internal carotid artery dissection. *Neurology* 2000;54:442.
- [32] Molina CA, Alvarez-Saban J, Schonewille W, et al. A Cerebral microembolism in acute spontaneous internal carotid artery dissection. *Neurology* 2000;55:1738–40.
- [33] Korchi AM, Cuvincini V, Caetano J, et al. Imaging of the cavernous sinus lesions. *Diagn Interv Imaging* 2014;95:849–59.
- [34] Charbonneau F, Williams M, Lafitte F, Hérán F. No more fear of the cavernous sinuses! *Diagn Interv Imaging* 2013;94:1003–16.
- [35] Thurtell MJ, Chiu AL, Goold LA, et al. Neuro-ophthalmology of invasive fungal sinusitis: 14 consecutive patients and a review of the literature. *Clin Experiment Ophthalmol* 2013;41:567–76.
- [36] Choi KD, Choi SY, Kim JS, et al. Acquired ocular motor nerve palsy in neurology clinics: a prospective multicenter study. *J Clin Neurol* 2019;15:221–7.