

## BIBLIOGRAPHIE

- [1] Kidd D. The optic chiasm. *Clin Anat* 2014;27:1149–58.
- [2] Toanen V, Héran F, Vignal C. Chiasma : anatomie, organisation fonctionnelle et neuro-imagerie. *Encycl Méd Chir (Elsevier, Paris). Ophtalmologie* 2017;14:17. 21-008-A-20.
- [3] Vignal-Clermont C, Tilikete C, Miléa D. *Neuro-ophtalmologie*. Paris : Elsevier-Masson 2016.
- [4] Won HS, Han SH, Oh CS, et al. Topographic variations of the optic chiasm and the foramen diaphragm sellae. *Surg Radiol Anat* 2010;32:653–7.
- [5] Tieger MG, Hedges TR, Ho J, et al. Ganglion cell complex loss in chiasmal compression by brain tumors. *J Neuroophthalmol* 2017;37:7–12.
- [6] Héran F, Lafitte F. *Imagerie en ophtalmologie pour les radiologues*. Issy-les-Moulineaux : Elsevier Masson 2018.
- [7] Chanson P, Raverot G, Castinetti F, et al. French Endocrinology Society non-functioning pituitary adenoma work-group. Management of clinically non-functioning pituitary adenoma. *Ann Endocrinol (Paris)* 2015;76:239–47.
- [8] Freda PU, Post KD. Differential diagnosis of sellar masses. *Endocrinol Metab Clin North Am* 1999;28:81–117.
- [9] Wildemberg LE, Glezer A, Bronstein MD, Gadelha MR. Apoplexy in nonfunctioning pituitary adenomas. *Pituitary* 2018;21:138–44.