

## BIBLIOGRAPHIE

- [1] Kidd D, Newman N, Biousse V. Neuro-Ophthalmology. Issy-les-Moulineaux: Elsevier Masson 2008.
- [2] Safran A, Vighetto A, Landis T, et al. Neuro-ophtalmologie. Rapport de la Société française d'ophtalmologie. Paris: Masson; 2004.
- [3] Schiefer U, Wilhelm H, Hart W. Clinical neuro-ophthalmology: a practical guide. Berlin-Heidelberg: Springer-Verlag 2007.
- [4] Prasad S. Diagnostic neuroimaging in neuro-ophthalmic disorders. Continuum (Minneap Minn) 2014;20:1023–62. 4 Neuro-ophthalmology.
- [5] Menjot de Champfleur N, Menjot de Champfleur S, Leboucq N, et al. Imagerie du chiasma optique et des voies optiques rétro-chiasmatiques. Journal de Radiologie Diagnostique et Interventionnelle 2013;94:964–78.

- [6] Brown HD, Woodall RL, Kitching RE, et al. Using magnetic resonance imaging to assess visual deficits: a review. Ophthalmic Physiol Opt 2016;36:240–65.
- [7] Rutland JW, Padormo F, Yim CK, et al. Quantitative assessment of secondary white matter injury in the visual pathway by pituitary adenomas: a multimodal study at 7-Tesla MRI. J Neurosurg 2019;1–10.
- [8] Hannoun S, Heidelberg D, Hourani R, et al. Diagnostic value of 3DFLAIR in clinical practice for the detection of infratentorial lesions in multiple sclerosis in regard to dual echo T2 sequences. Eur J Radiol 2018;102:146–51.