

BIBLIOGRAPHIE

Dystrophie de Fuchs (OMIM 136800)

- [1] Matthaehi M, Hribek A, Clahsen T, et al. Fuchs endothelial corneal dystrophy : clinical, genetic, pathophysiologic, and therapeutic aspects. *Annu Rev Vis Sci* 2019;5:151–75.
- [2] Borderie V, Baudrimont M, Bourcier T, et al. Cornea guttata et dystrophie de Fuchs. *J Fr Ophtalmol* 1999;22:987–1002.
- [3] Hogan MJ, Wood I, Fine M. Fuchs' endothelial dystrophy of the cornea. 29th Sanford Gifford Memorial lecture. *Am J Ophthalmol* 1974;78:363–83.
- [4] Lorenzetti DW, Uotila MH, Parikh N, Kaufman HE. Central cornea guttata. Incidence in the general population *Am J Ophthalmol* 1967;64:1155–8.
- [5] Weisenthal RW, Streeten BW. Posterior membrane dystrophies. In: Krachmer JH, Mannis MJ, Holland EJ, editors. *Cornea*. St. Louis: Mosby-Year Book; 1997. p. 1063–90.
- [6] Kaufman HE, Capella JA, Robbins JE. The human corneal endothelium. of Medicine, USA. *Am J Ophthalmol* 1966;61:835–41.
- [7] Borderie VM, Sabolic V, Touzeau O, et al. Screening human donor corneas during organ culture for the presence of guttae. *Br J Ophthalmol* 2001;85:272–6.
- [8] Zoega GM, Fujisawa A, Sasaki H, et al. Prevalence and risk factors for cornea guttata in the Reykjavik Eye Study. *Ophthalmology* 2006;113:565–9.
- [9] Higa A, Sakai H, Sawaguchi S, et al. Prevalence of and risk factors for cornea guttata in a population-based study in a southwestern island of Japan : the Kumejima study. *Arch Ophthalmol* 2011;129:332–6.
- [10] Zoega GM, Arnarsson A, Sasaki H, et al. The 7-year cumulative incidence of cornea guttata and morphological changes in the corneal endothelium in the Reykjavik Eye Study. *Acta Ophthalmol* 2013;91:212–8.
- [11] Mootha VV, Hansen B, Rong Z, et al. Fuchs' endothelial corneal dystrophy and RNA foci in patients with myotonic dystrophy. *Invest Ophthalmol Vis Sci* 2017;58:4579–85.
- [12] Olsen T. Is there an association between Fuchs' endothelial dystrophy and cardiovascular disease ? *Graefes Arch Clin Exp Ophthalmol* 1984;221:239–40.
- [13] Lipman RM, Rubenstein JB, Torczynski E. Keratoconus and Fuchs' corneal endothelial dystrophy in a patient and her family. *Arch Ophthalmol* 1990;108:993–4.
- [14] Nagarsheth M, Singh A, Schmotzer B, et al. Relationship between Fuchs endothelial corneal dystrophy severity and glaucoma and/or ocular hypertension. *Arch Ophthalmol* 2012;130:1384–8.
- [15] Pitts JF, Jay JL. The association of Fuchs's corneal endothelial dystrophy with axial hypermetropia, shallow anterior chamber, and angle closure glaucoma. *Br J Ophthalmol* 1990;74:601–4.
- [16] Rao GP, Kaye SB, Agius-Fernandez A. Central corneal endothelial guttae and age-related macular degeneration : is there an association ? *Indian J Ophthalmol* 1998;46:145–7.
- [17] Waring 3rd GO, Bourne WM, Edelhauser HF, Kenyon KR. The corneal endothelium. Normal and pathologic structure and function. *Ophthalmology* 1982;89:531–90.
- [18] Rodrigues MM, Krachmer JH, Hackett J, et al. Fuchs' corneal dystrophy. A clinicopathologic study of the variation in corneal edema. *Ophthalmology* 1986;93:789–96.
- [19] Kenney MC, Labermeier U, Hinds D, Waring GO 3rd. Characterization of the Descemet's membrane/posterior collagenous layer isolated from Fuchs' endothelial dystrophy corneas. *Exp Eye Res* 1984 39 : 267-77.
- [20] Gottsch JD, Zhang C, Sundin OH, et al. Fuchs corneal dystrophy: aberrant collagen distribution in an L450W mutant of the COL8A2 gene. *Invest Ophthalmol Vis Sci* 2005;46:4504–11.
- [21] Matthaehi M, Hu J, Kallay L, et al. Endothelial cell microRNA expression in human late-onset Fuchs' dystrophy. *Invest Ophthalmol Vis Sci* 2014;55:216–25.
- [22] Okumura N, Hashimoto K, Kitahara M, et al. Activation of TGF-beta signaling induces cell death via the unfolded protein response in Fuchs endothelial corneal dystrophy. *Sci Rep* 2017;7:6801–17.
- [23] Jurkunas UV, Bitar M, Rawe I. Colocalization of increased transforming growth factor-beta-induced protein (TGFB1p) and Clusterin in Fuchs endothelial corneal dystrophy. *Invest Ophthalmol Vis Sci* 2009;50:1129–36.
- [24] Poulsen ET, Dyrland TF, Runager K, et al. Proteomics of Fuchs' endothelial corneal dystrophy support that the extracellular matrix of Descemet's membrane is disordered. *J Proteome Res* 2014;13:4659–67.
- [25] Weller JM, Zenkel M, Schlotzer-Schrehardt U, et al. Extracellular matrix alterations in late-onset Fuchs' corneal dystrophy. *Invest Ophthalmol Vis Sci* 2014;55:3700–8.
- [26] Borderie VM, Baudrimont M, Vallee A, et al. Corneal endothelial cell apoptosis in patients with Fuchs' dystrophy. *Invest Ophthalmol Vis Sci* 2000;41:2501–5.
- [27] Li QJ, Ashraf MF, Shen DF, et al. The role of apoptosis in the pathogenesis of Fuchs endothelial dystrophy of the cornea. *Arch Ophthalmol* 2001;119:1597–604.
- [28] Adamis AP, Filatov V, Tripathi BJ, Tripathi RC. Fuchs' endothelial dystrophy of the cornea. *Surv Ophthalmol* 1993;38:149–68.
- [29] Iwamoto T, DeVoe AG. Electron microscopic studies on Fuchs' combined dystrophy. II. Anterior portion of the cornea. *Invest Ophthalmol* 1971;10:29–40.
- [30] McCartney MD, Wood TO, McLaughlin BJ. Immunohistochemical localization of ATPase in human dysfunctional corneal endothelium. *Curr Eye Res* 1987;6:1479–86.
- [31] McCartney MD, Wood TO, McLaughlin BJ. Moderate Fuchs' endothelial dystrophy ATPase pump site density. *Invest Ophthalmol Vis Sci* 1989;30:1560–4.
- [32] Tuberville AW, Wood TO, McLaughlin BJ. Cytochrome oxidase activity of Fuchs' endothelial dystrophy. *Curr Eye Res* 1986;5:939–47.
- [33] Mandell RB, Polse KA, Brand RJ, et al. Corneal hydration control in Fuchs' dystrophy. *Invest Ophthalmol Vis Sci* 1989;30:845–52.
- [34] Saini JS, Mittal S. In vivo quantification of corneal endothelium function. *Acta Ophthalmol Scand* 1996;74:468–72.
- [35] Wilson SE, Bourne WM, O'Brien PC, Brubaker RF. Endothelial function and aqueous humor flow rate in patients with Fuchs' dystrophy. *Am J Ophthalmol* 1988;106:270–8.
- [36] Burns RR, Bourne WM, Brubaker RF. Endothelial function in patients with cornea guttata. *Invest Ophthalmol Vis Sci* 1981;20:77–85.
- [37] Bourne WM. Clinical estimation of corneal endothelial pump function. *Trans Am Ophthalmol Soc* 1998;96:229–39.
- [38] Kocaba V, Katikireddy KR, Gipson I, et al. Association of the gutta-induced microenvironment with corneal endothelial cell behavior and demise in Fuchs endothelial corneal dystrophy. *JAMA Ophthalmol* 2018;136:886–92.
- [39] Raphael B, Lange T, Wood TO, McLaughlin BJ. Growth of human corneal endothelium on altered Descemet's membrane. *Cornea* 1992;11:242–9.
- [40] Bahn CF, Falls HF, Varley GA, et al. Classification of corneal endothelial disorders based on neural crest origin. *Ophthalmology* 1984;91:558–63.
- [41] Bourne WM, Johnson DH, Campbell RJ. The ultrastructure of Descemet's membrane. III. Fuchs' dystrophy. *Arch Ophthalmol* 1982;100:1952–5.
- [42] Rosenblum P, Stark WJ, Maumenee IH, et al. Hereditary Fuchs' dystrophy. *Am J Ophthalmol* 1980;90:455–62.
- [43] Krachmer JH, Purcell Jr JJ, Young CW, Bucher KD. Corneal endothelial dystrophy. A study of 64 families. *Arch Ophthalmol* 1978;96:2036–9.
- [44] Magovern M, Beauchamp GR, McTigue JW, et al. Inheritance of Fuchs' combined dystrophy. *Ophthalmology* 1979;86:1897–923.
- [45] Afshari NA, Pittard AB, Siddiqui A, Klintworth GK. Clinical study of Fuchs corneal endothelial dystrophy leading to penetrating keratoplasty: a 30-year experience. *Arch Ophthalmol* 2006;124:777–80.
- [46] Gottsch JD, Sundin OH, Liu SH, et al. Inheritance of a novel COL8A2 mutation defines a distinct early-onset subtype of fuchs corneal dystrophy. *Invest Ophthalmol Vis Sci* 2005;46:1934–9.
- [47] Mootha VV, Hansen B, Rong Z, et al. Fuchs' endothelial corneal dystrophy and RNA foci in patients with myotonic dystrophy. *Invest Ophthalmol Vis Sci* 2017;58:4579–85.
- [48] Matthaehi M, Zhu AY, Kallay L, et al. Transcript profile of cellular senescence-related genes in Fuchs endothelial corneal dystrophy. *Exp Eye Res* 2014;129:13–7.

- [49] Jurkunas UV, Bitar MS, Funaki T, Azizi B. Evidence of oxidative stress in the pathogenesis of fuchs endothelial corneal dystrophy. *Am J Pathol* 2010;177:2278–89.
- [50] Iwamoto T, DeVoe AG. Electron microscopic studies on Fuchs' combined dystrophy. I. Posterior portion of the cornea. *Invest Ophthalmol* 1971;10:9–28.
- [51] Amin SR, Baratz KH, McLaren JW, Patel SV. Corneal abnormalities early in the course of Fuchs' endothelial dystrophy. *Ophthalmology* 2014;121:2325–33.
- [52] Schrems-Hoesl LM, Schrems WA, Cruzat A, et al. Cellular and subbasal nerve alterations in early stage Fuchs' endothelial corneal dystrophy: an in vivo confocal microscopy study. *Eye (Lond)* 2013;27:42–9.
- [53] Hirano K, Klintworth GK, Zhan Q, et al. Beta ig-h3 is synthesized by corneal epithelium and perhaps endothelium in Fuchs' dystrophic corneas. *Curr Eye Res* 1996;15:965–72.
- [54] Calandra A, Chwa M, Kenney MC. Characterization of stroma from Fuchs' endothelial dystrophy corneas. *Cornea* 1989;8:90–7.
- [55] Muller LJ, Pels L, Vrensen GF. Novel aspects of the ultrastructural organization of human corneal keratocytes. *Invest Ophthalmol Vis Sci* 1995;36:2557–67.
- [56] Sasaki Y, Tuberville AW, Wood TO, McLaughlin BJ. Freeze fracture study of human corneal endothelial dysfunction. *Invest Ophthalmol Vis Sci* 1986;27:480–5.
- [57] Alexander RA, Grierson I, Garner A. Oxytalan fibers in Fuch's endothelial dystrophy. *Arch Ophthalmol* 1981;99:1622–7.
- [58] Levy SG, Moss J, Sawada H, et al. The composition of wide-spaced collagen in normal and diseased Descemet's membrane. *Curr Eye Res* 1996;15:45–52.
- [59] Davies Y, Fullwood NJ, Marcyniuk B, et al. Keratan sulphate in the trabecular meshwork and cornea. *Curr Eye Res* 1997;16:677–86.
- [60] Polack FM. The posterior corneal surface in Fuchs' dystrophy. Scanning electron microscope study. *Invest Ophthalmol* 1974;13:913–22.
- [61] Carden RG. Optometric management of Fuchs's endothelial dystrophy. *Am J Optom Physiol Opt* 1978;55:642–6.
- [62] Wilson SE, Bourne WM. Fuchs' dystrophy. *Cornea* 1988;7:2–18.
- [63] Watanabe S, Oie Y, Fujimoto H, et al. Relationship between corneal guttae and quality of vision in patients with mild Fuchs' endothelial corneal dystrophy. *Ophthalmology* 2015;122:2103–9.
- [64] Wacker K, Baratz KH, Bourne WM, Patel SV. Patient-reported visual disability in Fuchs' endothelial corneal dystrophy measured by the visual function and corneal health status instrument. *Ophthalmology* 2018;125:1854–61.
- [65] Borderie V. *Imagerie de la cornée et de la surface oculaire*. Paris: Éditions Med'com; 2014.
- [66] Shousha MA, Perez VL, Wang J, et al. Use of ultra-high-resolution optical coherence tomography to detect in vivo characteristics of Descemet's membrane in Fuchs' dystrophy. *Ophthalmology* 2010;117:1220–7.
- [67] Repp DJ, Hodge DO, Baratz KH, et al. Fuchs' endothelial corneal dystrophy: subjective grading versus objective grading based on the central-to-peripheral thickness ratio. *Ophthalmology* 2013;120:687–94.
- [68] Ghoulali W, Grieve K, Bellefqî S, et al. Full-field optical coherence tomography of human donor and pathological corneas. *Curr Eye Res* 2015;40:526–34.
- [69] Schnitzer JI, Krachmer JH. A specular microscopic study of families with endothelial dystrophy. *Br J Ophthalmol* 1981;65:396–400.
- [70] Laing RA, Leibowitz HM, Oak SS, et al. Endothelial mosaic in Fuchs' dystrophy. A qualitative evaluation with the specular microscope. *Arch Ophthalmol* 1981;99:80–3.
- [71] Loewenstein A, Hourvitz D, Goldstein M, et al. Association of fuchs' corneal endothelial dystrophy with angle-closure glaucoma. *J Glaucoma* 1994;3:201–5.
- [72] Brooks AM, Grant G, Gillies WE. The significance of anterior chamber depth in Fuchs' corneal dystrophy and cornea guttata. *Cornea* 1994;13:131–5.
- [73] Mustonen RK, McDonald MB, Srivannaboon S, et al. In vivo confocal microscopy of Fuchs' endothelial dystrophy. *Cornea* 1998;17:493–503.
- [74] Kaufman SC, Beuerman RW, Kaufman HE. Diagnosis of advanced Fuchs' endothelial dystrophy with the confocal microscope. *Am J Ophthalmol* 1993;116:652–3.
- [75] Brooks AM, Grant G, Gillies WE. The use of specular microscopy to investigate unusual findings in the corneal endothelium and its adjacent structures. *Aust N Z J Ophthalmol* 1988;16:235–43.

Caractéristiques génétiques de la dystrophie endothéliale cornéenne de Fuchs

- [76] Matthaëi M, Hribek A, Clahsen T, et al. Fuchs Endothelial corneal dystrophy: clinical, genetic, pathophysiologic, and therapeutic aspects. *Ann Rev Vision Sci* 2019;5:151–75.
- [77] Mehta JS, Vithana EN, Tan DT, et al. Analysis of the posterior polymorphous corneal dystrophy 3 gene, TCF8, in late-onset Fuchs endothelial corneal dystrophy. *Invest Ophthalmol Vis Sci* 2008;49:184–8.
- [78] Soh YQ, Kocaba V, Pinto M, Mehta JS. Fuchs endothelial corneal dystrophy and corneal endothelial diseases : East meets West. *Eye (Lond)* 2020;34:427–41.
- [79] Matthaëi M, Sandhaeger H, Hermel M, et al. Changing indications in penetrating keratoplasty: a systematic review of 34 years of global reporting. *Transplantation* 2017;101:1387–99.
- [80] Zoega GM, Fujisawa A, Sasaki H, et al. Prevalence and risk factors for cornea guttata in the Reykjavik Eye Study. *Ophthalmology* 2006;113:565–9.
- [81] Lorenzetti DW, Uotila MH, Parikh N, Kaufman HE. Central cornea guttata. Incidence in the general population. *Am J Ophthalmol* 1967;64:1155–8.
- [82] Higa A, Sakai H, Sawaguchi S, et al. Prevalence of and risk factors for cornea guttata in a population-based study in a southwestern island of Japan : the Kumejima study. *Arch Ophthalmol* 2011;129:332–6.
- [83] Gottsch JD, Sundin OH, Liu SH, et al. Inheritance of a novel COL8A2 mutation defines a distinct early-onset subtype of fuchs corneal dystrophy. *Invest Ophthalmol Vis Sci* 2005;46:1934–9.
- [84] Krachmer JH, Purcell Jr JJ, Young CW, Bucher KD. Corneal endothelial dystrophy. A study of 64 families. *Arch Ophthalmol* 1978;96:2036–9.
- [85] Afshari NA, Pittard AB, Siddiqui A, Klintworth GK. Clinical study of Fuchs corneal endothelial dystrophy leading to penetrating keratoplasty: a 30-year experience. *Arch Ophthalmol* 2006;124:777–80.
- [86] Biswas S, Munier FL, Yardley J, et al. Missense mutations in COL8A2, the gene encoding the alpha2 chain of type VIII collagen, cause two forms of corneal endothelial dystrophy. *Hum Mol Genet* 2001;10:2415–23.
- [87] Mok JW, Kim HS, Joo CK. Q455V mutation in COL8A2 is associated with Fuchs' corneal dystrophy in Korean patients. *Eye (Lond)* 2009;23:895–903.
- [88] Magovern M, Beauchamp GR, McTigue JW, et al. Inheritance of Fuchs' combined dystrophy. *Ophthalmology* 1979;86:1897–923.
- [89] Liskova P, Prescott Q, Bhattacharya SS, Tuft SJ. British family with early-onset Fuchs' endothelial corneal dystrophy associated with p.L450W mutation in the COL8A2 gene. *Br J Ophthalmol* 2007;91:1717–8.
- [90] Aldave AJ, Rayner SA, Salem AK, et al. No pathogenic mutations identified in the COL8A1 and COL8A2 genes in familial Fuchs corneal dystrophy. *Invest Ophthalmol Vis Sci* 2006;47:3787–90.
- [91] Gupta R, Kumawat BL, Paliwal P, et al. Association of ZEB1 and TCF4 rs613872 changes with late onset Fuchs endothelial corneal dystrophy in patients from northern India. *Mol Vis* 2015;21:1252–60.
- [92] Jun AS, Meng H, Ramanan N, et al. An alpha 2 collagen VIII transgenic knock-in mouse model of Fuchs endothelial corneal dystrophy shows early endothelial cell unfolded protein response and apoptosis. *Hum Mol Genet* 2012;21:384–93.
- [93] Meng H, Matthaëi M, Ramanan N, et al. L450W and Q455K Col8a2 knock-in mouse models of Fuchs endothelial corneal dystrophy show distinct phenotypes and evidence for altered autophagy. *Invest Ophthalmol Vis Sci* 2013;54:1887–97.
- [94] Engler C, Kelliher C, Spitze AR, et al. Unfolded protein response in Fuchs endothelial corneal dystrophy: a unifying pathogenic pathway? *Am J Ophthalmol* 2010;149:194–202.e2.

- [95] Vedana G, Villarreal Jr G, Jun AS. Fuchs endothelial corneal dystrophy: current perspectives. *Clin Ophthalmol* 2016;10:321–30.
- [96] Amiel J, Rio M, de Pontual L, et al. Mutations in TCF4, encoding a class I basic helix-loop-helix transcription factor, are responsible for Pitt-Hopkins syndrome, a severe epileptic encephalopathy associated with autonomic dysfunction. *Am J Hum Genet* 2007;80:988–93.
- [97] Navarrete K, Pedroso I, De Jong S, et al. TCF4 (e2-2 ITF2) : a schizophrenia-associated gene with pleiotropic effects on human disease. *Am J Med Genet B Neuropsychiatr Genet* 2013;162:1–16.
- [98] Forrest MP, Waite AJ, Martin-Rendon E, Blake DJ. Knockdown of human TCF4 affects multiple signaling pathways involved in cell survival, epithelial to mesenchymal transition and neuronal differentiation. *PLoS one* 2013;8, e73169.
- [99] Baratz KH, Tosakulwong N, Ryu E, et al. E2-2 protein and Fuchs's corneal dystrophy. *N Engl J Med* 2010;363:1016–24.
- [100] Wieben ED, Aleff RA, Tosakulwong N, et al. A common trinucleotide repeat expansion within the transcription factor 4 (TCF4, E2-2) gene predicts Fuchs corneal dystrophy. *PLoS One* 2012;7.
- [101] Zhang J, McGhee CNJ, Patel DV. The molecular basis of Fuchs' endothelial corneal dystrophy. *Mol Diag Ther* 2019;23:97–112.
- [102] Mootha VV, Gong X, Ku HC, Xing C. Association and familial segregation of CTG18.1 trinucleotide repeat expansion of TCF4 gene in Fuchs' endothelial corneal dystrophy. *Invest Ophthalmol Vis Sci* 2014;55:33–42.
- [103] Wieben ED, Aleff RA, Eckloff BW, et al. Comprehensive assessment of genetic variants within TCF4 in Fuchs' endothelial corneal dystrophy. *Invest Ophthalmol Vis Sci* 2014;55:6101–7.
- [104] Nanda GG, Padhy B, Samal S, et al. Genetic association of TCF4 intronic polymorphisms, CTG18.1 and rs17089887, with Fuchs' endothelial corneal dystrophy in an Indian population. *Invest Ophthalmol Vis Sci* 2014;55:7674–80.
- [105] Xing C, Gong X, Hussain I, et al. Transethnic replication of association of CTG18.1 repeat expansion of TCF4 gene with Fuchs' corneal dystrophy in Chinese implies common causal variant. *Invest Ophthalmol Vis Sci* 2014;55:7073–8.
- [106] Vasanth S, Eghrari AO, Gapsis BC, et al. Expansion of CTG18.1 trinucleotide repeat in TCF4 is a potent driver of Fuchs' corneal dystrophy. *Invest Ophthalmol Vis Sci* 2015;56:4531–6.
- [107] Nakano M, Okumura N, Nakagawa H, et al. Trinucleotide repeat expansion in the TCF4 gene in Fuchs' endothelial corneal dystrophy in Japanese. *Invest Ophthalmol Vis Sci* 2015;56:4865–9.
- [108] Rao BS, Tharigopala A, Rachapalli SR, et al. Association of polymorphisms in the intron of TCF4 gene to late-onset Fuchs endothelial corneal dystrophy : An Indian cohort study. *Indian J Ophthalmol* 2017;65:931–5.
- [109] Eghrari AO, Vasanth S, Wang J, et al. CTG18.1 Expansion in TCF4 increases likelihood of transplantation in Fuchs corneal dystrophy. *Cornea* 2017;36:40–3.
- [110] Foja S, Luther M, Hoffmann K, et al. CTG18.1 repeat expansion may reduce TCF4 gene expression in corneal endothelial cells of German patients with Fuchs' dystrophy. *Graefes Arch Clin Exp Ophthalmol* 2017;255:1621–31.
- [111] Kuot A, Hewitt AW, Snibson GR, et al. TGC repeat expansion in the TCF4 gene increases the risk of Fuchs' endothelial corneal dystrophy in Australian cases. *PLoS One* 2017;12.
- [112] Zarouchlioti C, Sanchez-Pintado B, Hafford Tear NJ, et al. Antisense therapy for a common corneal dystrophy ameliorates TCF4 repeat expansion-mediated toxicity. *Am J Hum Genet* 2018;102:528–39.
- [113] Zhang J, Patel DV. The pathophysiology of Fuchs' endothelial dystrophy - a review of molecular and cellular insights. *Exp Eye Res* 2015;130:97–105.
- [114] Soliman AZ, Xing C, Radwan SH, et al. Correlation of severity of Fuchs endothelial corneal dystrophy with triplet repeat expansion in TCF4. *JAMA Ophthalmol* 2015;133:1386–91.
- [115] Iliff BW, Riazuddin SA, Gottsch JD. The genetics of Fuchs' corneal dystrophy. *Expert Rev Ophthalmol* 2012;7:363–75.
- [116] Riazuddin SA, McGlumphy EJ, Yeo WS, et al. Replication of the TCF4 intronic variant in late-onset Fuchs corneal dystrophy and evidence of independence from the FCD2 locus. *Invest Ophthalmol Vis Sci* 2011;52:2825–9.
- [117] Stamler JF, Roos BR, Wagoner MD, et al. Confirmation of the association between the TCF4 risk allele and Fuchs endothelial corneal dystrophy in patients from the Midwestern United States. *Ophthalmic Genet* 2013;34:32–4.
- [118] Oldak M, Ruszkowska E, Udziela M, et al. Fuchs endothelial corneal dystrophy : strong association with rs613872 not paralleled by changes in corneal endothelial TCF4 mRNA level. *BioMed Res Int* 2015;2015.
- [119] Afshari NA, Igo Jr RP, Morris NJ, et al. Genome-wide association study identifies three novel loci in Fuchs endothelial corneal dystrophy. *Nat Commun* 2017;8.
- [120] Eghrari AO, McGlumphy EJ, Iliff BW, et al. Prevalence and severity of Fuchs corneal dystrophy in Tangier Island. *Am J Ophthalmol* 2012;153:1067–72.
- [121] Li D, Peng X, Sun H. Association of TCF4 polymorphisms and Fuchs' endothelial dystrophy : a meta-analysis. *BMC Ophthalmol* 2015;15.
- [122] Wang KJ, Jhanji V, Chen J, et al. Association of transcription factor 4 (TCF4) and protein tyrosine phosphatase, receptor type G (PTPRG) with corneal dystrophies in southern Chinese. *Ophthalmic Genet* 2014;35:138–41.
- [123] Thalamuthu A, Khor CC, Venkataraman D, et al. Association of TCF4 gene polymorphisms with Fuchs' corneal dystrophy in the Chinese. *Invest Ophthalmol Vis Sci* 2011;52:5573–8.
- [124] Minear MA, Li YJ, Rimmler J, et al. Genetic screen of African Americans with Fuchs endothelial corneal dystrophy. *Mol Vis* 2013;19:2508–16.
- [125] Du J, Aleff RA, Soragni E, et al. RNA toxicity and missplicing in the common eye disease Fuchs endothelial corneal dystrophy. *J Biol Chem* 2015;290:5979–90.
- [126] Hu J, Rong Z, Gong X, et al. Oligonucleotides targeting TCF4 triplet repeat expansion inhibit RNA foci and mis-splicing in Fuchs' dystrophy. *Hum Mol Genet* 2018;27:1015–26.
- [127] Krafchak CM, Pawar H, Moroi SE, et al. Mutations in TCF8 cause posterior polymorphous corneal dystrophy and ectopic expression of COL4A3 by corneal endothelial cells. *Am J Hum Genet* 2005;77:694–708.
- [128] Aldave AJ, Yellore VS, Yu F, et al. Posterior polymorphous corneal dystrophy is associated with TCF8 gene mutations and abdominal hernia. *Am J Med Genet A* 2007;143A:2549–56.
- [129] Riazuddin SA, Zaghoul NA, Al-Saif A, et al. Missense mutations in TCF8 cause late-onset Fuchs corneal dystrophy and interact with FCD4 on chromosome 9p. *Am J Hum Genet* 2010;86:45–53.
- [130] Vithana EN, Morgan PE, Ramprasad V, et al. SLC4A11 mutations in Fuchs endothelial corneal dystrophy. *Hum Mol Genet* 2008;17:656–66.
- [131] Vithana EN, Morgan P, Sundaresan P, et al. Mutations in sodium-borate cotransporter SLC4A11 cause recessive congenital hereditary endothelial dystrophy (CHED2). *Nat Genet* 2006;38:755–7.
- [132] Desir J, Moya G, Reish O, et al. Borate transporter SLC4A11 mutations cause both Harboyan syndrome and non-syndromic corneal endothelial dystrophy. *J Med Genet* 2007;44:322–6.
- [133] Malhotra D, Loganathan SK, Chiu AM, et al. Human corneal expression of SLC4A11, a gene mutated in endothelial corneal dystrophies. *Sci Rep* 2019;9.
- [134] Gottsch JD, Bowers AL, Margulies EH, et al. Serial analysis of gene expression in the corneal endothelium of Fuchs' dystrophy. *Invest Ophthalmol Vis Sci* 2003;44:594–9.
- [135] Riazuddin SA, Vithana EN, Seet LF, et al. Missense mutations in the sodium borate cotransporter SLC4A11 cause late-onset Fuchs corneal dystrophy. *Hum Mutat* 2010;31:1261–8.
- [136] Soumitra N, Loganathan SK, Madhavan D, et al. Biosynthetic and functional defects in newly identified SLC4A11 mutants and absence of COL8A2 mutations in Fuchs endothelial corneal dystrophy. *J Hum Genet* 2014;59:444–53.
- [137] Vilas GL, Loganathan SK, Quon A, et al. Oligomerization of SLC4A11 protein and the severity of FECD and CHED2 corneal dystrophies caused by SLC4A11 mutations. *Hum Mutat* 2012;33:419–28.
- [138] Riazuddin SA, Vasanth S, Katsanis N, Gottsch JD. Mutations in AGBL1 cause dominant late-onset Fuchs corneal dystrophy and alter protein-protein interaction with TCF4. *Am J Hum Genet* 2013;93:758–64.

- [139] Riazuddin SA, Parker DS, McGlumphy EJ, et al. Mutations in LOXHD1, a recessive-deafness locus, cause dominant late-onset Fuchs corneal dystrophy. *Am J Hum Genet* 2012;90:533–9.
- [140] Mori K, Moteki H, Kobayashi Y, et al. Mutations in LOXHD1 gene cause various types and severities of hearing loss. *Ann Otol Rhinol Laryngol* 2015;124(Suppl 1):135S–41S.
- [141] Jurkunas UV, Bitar MS, Rawe I, et al. Increased clusterin expression in Fuchs' endothelial dystrophy. *Invest Ophthalmol Vis Sci* 2008;49:2946–55.
- [142] Jurkunas UV, Bitar M, Rawe I. Colocalization of increased transforming growth factor-beta-induced protein (TGFBIP) and Clusterin in Fuchs endothelial corneal dystrophy. *Invest Ophthalmol Vis Sci* 2009;50:1129–36.
- [143] Weller JM, Zenkel M, Schlotzer-Schrehardt U, et al. Extracellular matrix alterations in late-onset Fuchs' corneal dystrophy. *Invest Ophthalmol Vis Sci* 2014;55:3700–8.
- [144] Kuot A, Hewitt AW, Griggs K, et al. Association of TCF4 and CLU polymorphisms with Fuchs' endothelial dystrophy and implication of CLU and TGFBIP proteins in the disease process. *Eur J Hum Genet* 2012;20(6):632–8.
- [145] Synowiec E, Wojcik KA, Izdebska J, et al. Polymorphisms of the homologous recombination gene RAD51 in keratoconus and Fuchs endothelial corneal dystrophy. *Dis Markers* 2013;35:353–62.
- [146] Wojcik KA, Synowiec E, Polakowski P, et al. Polymorphism of the flap endonuclease 1 gene in keratoconus and Fuchs endothelial corneal dystrophy. *Int J Mol Sci* 2014;15:14786–802.
- [147] Synowiec E, Wojcik KA, Izdebska J, et al. Polymorphisms of the apoptosis-related FAS and FAS ligand genes in keratoconus and Fuchs endothelial corneal dystrophy. *Tohoku J Exp Med* 2014;234:17–27.
- [148] Wojcik KA, Synowiec E, Polakowski P, et al. Variation in DNA base excision repair genes in Fuchs endothelial corneal dystrophy. *Med Sci Monit* 2015;21:2809–27.
- [149] Nanda GG, Kumar MV, Pradhan L, et al. rs4246215 is targeted by hsa-miR1236 to regulate FEN1 expression but is not associated with Fuchs' endothelial corneal dystrophy. *PloS One* 2018;13.
- [150] Matthaei M, Hu J, Kallay L, et al. Endothelial cell microRNA expression in human late-onset Fuchs' dystrophy. *Invest Ophthalmol Vis Sci* 2014;55:216–25.
- [151] Sundin OH, Jun AS, Broman KW, et al. Linkage of late-onset Fuchs corneal dystrophy to a novel locus at 13pTel-13q12.13. *Invest Ophthalmol Vis Sci* 2006;47:140–5.
- [152] Meadows DN, Eghrari AO, Riazuddin SA, et al. Progression of Fuchs corneal dystrophy in a family linked to the FCD1 locus. *Invest Ophthalmol Vis Sci* 2009;50:5662–6.
- [153] Sundin OH, Broman KW, Chang HH, et al. A common locus for late-onset Fuchs corneal dystrophy maps to 18q21.2-q21.32. *Invest Ophthalmol Vis Sci* 2006;47:3919–26.
- [154] Riazuddin SA, Eghrari AO, Al-Saif A, et al. Linkage of a mild late-onset phenotype of Fuchs corneal dystrophy to a novel locus at 5q33.1-q35.2. *Invest Ophthalmol Vis Sci* 2009;50:5667–71.
- [155] Riazuddin SA, Zaghoul NA, Al-Saif A, et al. Missense mutations in TCF8 cause late-onset Fuchs corneal dystrophy and interact with FCD4 on chromosome 9p. *Am J Hum Genet* 2010;86:45–53.
- [156] Afshari NA, Li YJ, Pericak-Vance MA, et al. Genome-wide linkage scan in Fuchs endothelial corneal dystrophy. *Invest Ophthalmol Vis Sci* 2009;50:1093–7.
- [157] Gain P, Jullienne R, He Z, et al. Global survey of corneal transplantation and eye banking. *JAMA Ophthalmol* 2016;134:167–73.
- [158] Louttit MD, Kopplin LJ, Igo Jr RP, et al. A multicenter study to map genes for Fuchs endothelial corneal dystrophy : baseline characteristics and heritability. *Cornea* 2012;31:26–35.
- [159] Flicek P, Ahmed I, Amode MR, et al. Ensembl 2013. *Nucleic Acids Res* 2013;41. D48–55.