Bibliographie

[1] Ashena Z, Gallagher S, Naveed H, et al. Comparison of anterior corneal aberrometry, keratometry and pupil size with scheimpflug tomography and ray tracing aberrometer. Vision (Basel) 2022;6(1):18.
[2] Wang Y, Zhao K, Jin Y, et al. Changes of higher order aberration with various pupil sizes in the myopic eye. Pt 2J. Refract Surg 2003;19:S270–4.
[3] Wang L, Dai E, Koch DD, Nathoo A. Optical aberrations of the human anterior cornea. J Cataract Refract Surg 2003;29:1514–21.
[4] Atchison DA, Suheimat M, Mathur A, et al. Anterior corneal, posterior corneal, and lenticular contributions to ocular aberrations. Investig Ophthalmol Vis Sci 2016;57:5263–70.
[5] Hwang JS, Lee YP, Bae SH, et al. Utility of the optical quality analysis system for decision-making in cataract surgery. BMC Ophthalmol 2018;18(1):231.
[6] Miażdżyk M, Consejo A, Iskander DR. OCT based corneal densitometry : the confounding effect of epithelial speckle. Biomed Opt Express 2023;14(8):3871–80.
[7] Liang J, Williams DR. Aberrations and retinal image quality of the normal human eye. J Opt Soc Am A Opt Image Sci Vis 1997;14(11):2873–83.