Bibliographie

[1] Lizzi FL, Burt WJ, Coleman DJ. Effects of ocular structures on propagation of ultrasound in the eye. Arch Ophthalmol 1970;84(5):635–40.
[2] Reinstein DZ, Archer TJ, Gobbe M, et al. Repeatability of layered corneal pachymetry with the Artemis very high-frequency digital ultrasound arc-scanner. J Refract Surg 2010;26(9):646–59.
[3] Izatt JA, Hee MR, Swanson EA, et al. Micrometer-scale resolution imaging of the anterior eye in vivo with optical coherence tomography. Arch Ophthalmol 1994;112:1584–9.
[4] Shankar H, Taranath D, Santhirathelagan CT, et al. Anterior segment biometry with the Pentacam : comprehensive assessment of repeatability of automated measurements. J Cataract Refract Surg 2008;34(1):103–13.
[5] Rozema JJ, Wouters K, Mathysen DG, et al. Overview of the repeatability, reproducibility, and agreement of the biometry values provided by various ophthalmic devices. Am J Ophthalmol 2014;158(6):1111–20.
[6] Doughty MJ, Zaman ML. Human corneal thickness and its impact on intraocular pressure measures : a review and meta-analysis approach. Surv Ophthalmol 2000;44:367–408.
[7] La Rosa FA, Gross RL, Orengo-Nania S. Central corneal thickness of Caucasians and African Americans in glaucomatous and non glaucomatous populations. Arch Ophthalmol 2001;119(1):23–7.