Bibliographie

[1] Altamirano F, Ortiz-Morales G, O'Connor-Cordova MA, et al. Fuchs endothelial corneal dystrophy : an updated review. Int Ophthalmol 2024;44. 61-24.
[2] Weiss JS, Rapuano CJ, Seitz B, et al. IC3D classification of corneal dystrophies-Edition 3. Cornea 2024;43:466–527.
[3] Ong Tone S, Kocaba V, Böhm M, et al. Fuchs endothelial corneal dystrophy : the vicious cycle of Fuchs pathogenesis. Prog Retin Eye Res 2021 Jan;80, 100863.
[4] Evans CJ, Dudakova L, Skalicka P, et al. Schnyder corneal dystrophy and associated phenotypes caused by novel and recurrent mutations in the UBIAD1 gene. BMC Ophthalmol 2018;18. 250-18.
[5] Henriquez-Recine MAH, Lima KSM, Garcia EV, et al. Heredity and in vivo confocal microscopy of punctiform and polychromatic pre-Descemet dystrophy. Graefes Arch Clin Exp Ophthalmol 2018;256(9):1661–7.
[6] Patel SP, Parker MD. SLC4A11 and the pathophysiology of congenital hereditary endothelial dystrophy. Biomed Res Int 2015;2015, 475392.
[7] Kim MJ, Frausto RF, Rosenwasser GO, et al. Posterior amorphous corneal dystrophy is associated with a deletion of small leucine-rich proteoglycans on chromosome 12. PloS One 2014;9(4), e95037.
[8] Nowinska AK, Wylegala E, Teper S, et al. Phenotype-genotype correlation in patients with Schnyder corneal dystrophy. Cornea 2014;33:497–503.
[9] Nowinska AK, Wylegala E, Teper S, et al. Phenotype and genotype analysis in patients with macular corneal dystrophy. Br J Ophthalmol 2014;98:1514–21.
[10] Louttit MD, Kopplin LJ, Igo Jr RP, et al. A multicenter study to map genes for Fuchs endothelial corneal dystrophy : baseline characteristics and heritability. Cornea 2012;31:26–35.
[11] Paliwal P, Sharma A, Tandon R, et al. TGFBI mutation screening and genotype-phenotype correlation in north Indian patients with corneal dystrophies. Mol Vis 2010;16:1429–38.
[12] Kobayashi A, Sugiyama K. In vivo laser confocal microscopy findings for Bowman's layer dystrophies (Thiel-Behnke and Reis-Bucklers corneal dystrophies). Ophthalmology 2007;114:69–75.
[13] Magovern M, Beauchamp GR, McTigue JW, et al. Inheritance of Fuchs' combined dystrophy. Ophthalmology 1979;86:1897–923.
[14] Biswas S, Munier FL, Yardley J, et al. Missense mutations in COL8A2, the gene encoding the alpha2 chain of type VIII collagen, cause two forms of corneal endothelial dystrophy. Hum Mol Genet 2001;10:2415–23.
[15] Klintworth GK. The molecular genetics of the corneal dystrophies- -current status. Front Biosci 2003;8:687–713.
[16] Gottsch JD, Sundin OH, Liu SH, et al. Inheritance of a novel COL8A2 mutation defines a distinct early-onset subtype of Fuchs corneal dystrophy. Invest Ophthalmol Vis Sci 2005;46:1934–9.
[17] Weiss JS, Kruth HS, Kuivaniemi H, et al. Mutations in the UBIAD1 gene on chromosome short arm 1, region 36, cause Schnyder crystalline corneal dystrophy. Invest Ophthalmol Vis Sci 2007;48:5007–12.
[18] Orr A, Dube MP, Marcadier J, et al. Mutations in the UBIAD1 gene, encoding a potential prenyltransferase, are causal for Schnyder crystalline corneal dystrophy. PloS One 2007;2(8), e685.
[19] Chao-Shern C, DeDionisio LA, Jang JH, et al. Evaluation of TGFBI corneal dystrophy and molecular diagnostic testing. Eye (Lond) 2019;33:874–81.
[20] Yang GN, Sun YBY, Roberts PK, et al. Exploring single-cell RNA sequencing as a decision-making tool in the clinical management of Fuchs' endothelial corneal dystrophy. Prog Retin Eye Res 2024 Sep;102, 101286.