Bibliographie

[1] Eiberg H, Moller HU, Berendt I, Mohr J. Assignment of granular corneal dystrophy Groenouw type I (CDGG1) to chromosome 5q. Eur J Hum Genet 1994;2(2):132–8.
[2] Stone EM, Mathers WD, Rosenwasser GO, et al. Three autosomal dominant corneal dystrophies map to chromosome 5q. Nat Genet 1994;6(1):47–51.
[3] Munier FL, Korvatska E, Djemai A, et al. Kerato-epithelin mutations in four 5q31-linked corneal dystrophies. Nat Genet 1997;15(3):247–51.
[4] Niel-Butschi F, Kantelip B, Iwaszkiewicz J, et al. Genotype-phenotype correlations of TGFBI p.Leu509Pro, p.Leu509Arg, p.Val-613Gly, and the allelic association of p.Met502Val-p.Arg555Gln mutations. Mol Vis 2011;17:1192–202.
[5] Ellies P, Bejjani RA, Bourges JL, et al. Phototherapeutic keratectomy for BIGH3-linked corneal dystrophy recurring after penetrating keratoplasty. Ophthalmology 2003;110(6):1119–25.
[6] Behnke H, Thiel H. A previously unrecognized form of subepithelial corneal dystrophy inherited as an autosomal dominant. German Medical Monthly 1967;12(6):285.
[7] Küchle M, Green WR, Völcker HE, Barraquer J. Reevaluation of corneal dystrophies of Bowman's layer and the anterior stroma (Reis-Bücklers and Thiel-Behnke types) : a light and electron microscopic study of eight corneas and a review of the literature. Cornea 1995;14(4):333–54.
[8] Lisch W, Seitz B, Society SKotGO. Lattice corneal dystrophy, type 1 (LCD1) : an epithelial or stromal entity ? Invest Ophthalmol Vis Sci 2013;54(15). 4734-.
[9] Stock EL, Feder RS, O'Grady RB, et al. Lattice corneal dystrophy type IIIA : clinical and histopathologic correlations. Arch Ophthalmol 1991;109(3):354–8.
[10] Tsujikawa M, Kurahashi H, Tanaka T, et al. Identification of the gene responsible for gelatinous drop-like corneal dystrophy. Nat Genet 1999;21(4):420–3.
[11] Ren Z, Lin PY, Klintworth GK, et al. Allelic and locus heterogeneity in autosomal recessive gelatinous drop-like corneal dystrophy. Hum Genet 2002;110(6):568–77.
[12] Fujita S, Sameshima M, Hirashima S, Nakao K. Light and electron microscopic study of gelatinous drop-like corneal dystrophy with deeper stromal involvement. Nippon Ganka Gakkai Zasshi 1988;92(11):1744–57.
[13] Maeno S, Soma T, Tsujikawa M, et al. Efficacy of therapeutic soft contact lens in the management of gelatinous drop-like corneal dystrophy. Br J Ophthalmol 2020;104(2):241–6.