Bibliographie

[1] Tamimi A, Sheikhzadeh F, Ezabadi SG, et al. Post-LASIK dry eye disease : a comprehensive review of management and current treatment options. Front Med (Lausanne) 2023;10:1057685.

[2] Lee JB, Ryu CH, Kim J, et al. Comparison of tear secretion and tear film instability after photorefractive keratectomy and laser in situ keratomileusis. J Cataract Refract Surg 2000;26:1326–31.

[3] Yu EY, Leung A, Rao S, Lam DS. Effect of laser in situ keratomileusis on tear stability. Ophthalmology 2000;107:2131–5.

[4] Levinson BA, Rapuano CJ, Cohen EJ, et al. Referrals to the wills eye institute cornea service after laser in situ keratomileusis : reasons for patient dissatisfaction. J Cataract Refract Surg 2008;34:32–9.

[5] Bower KS, Sia RK, Ryan DS, et al. Chronic dry eye in photorefractive keratectomy and laser in situ keratomileusis : manifestations, incidence, and predictive factors. J Cataract Refract Surg 2015;41:2624–34.

[6] Vázquez A, Martínez-Plaza E, Fernández I, et al. Phenotypic characterization of patients developing chronic dry eye and pain after refractive surgery : a cross-sectional study. Ocul Surf 2022;26:63–74.

[7] Moshirfar M, Bhavsar UM, Durnford KM, et al. Neuropathic corneal pain following LASIK surgery : a retrospective case series. Ophthalmol Ther 2021;10:677–89.

[8] Albietz JM, Lenton LM. Management of the ocular surface and tear film before, during, and after laser in situ keratomileusis. J Refract Surg 2004;20:62–71.

[9] Shoja MR, Besharati MR. Dry eye after LASIK for myopia : incidence and risk factors. Eur J Ophthalmol 2007;17:1–6.

[10] Farahi A, Hashemi H, Mehravaran S, et al. Tear function evaluation in candidates of corneal laser refractive surgery for myopia. Eye Contact Lens 2014;40:91–4.

[11] Li Y, Li S, Zhou J, et al. Relationship between lipid layer thickness, incomplete blinking rate and tear film instability in patients with different myopia degrees after small-incision lenticule extraction. PLoS One 2020;15, e0230119.

[12] Moshirfar M, Fuhriman DA, Ali A, et al. Inflammatory bowel disease guidelines for corneal refractive surgery evaluation. J Clin Med 2022;11.

[13] Moshirfar M, Brown AH, Sulit CA, et al. Corneal refractive surgery considerations in patients with cystic fibrosis and cystic fibrosis transmembrane conductance regulator-related disorders. Int Med Case Rep J 2022;15:647–56.

[14] Toda I, Asano-Kato N, Hori-Komai Y, Tsubota K. Laser-assisted in situ keratomileusis for patients with dry eye. Arch Ophthalmol 2002;120:1024–8.

[15] Miles S, McGlathery W, Abernathie B. The importance of screening for laser-assisted in situ keratomileusis operation (LASIK) before prescribing isotretinoin. J Am Acad Dermatol 2006;54:180–1.

[16] Palme C, Mulrine F, McNeely RN, et al. Assessment of the correlation of the tear breakup time with quality of vision and dry eye symptoms after SMILE surgery. Int Ophthalmol 2022;42:1013–20.

[17] Albietz JM, Lenton LM, McLennan SG. Chronic dry eye and regression after laser in situ keratomileusis for myopia. J Cataract Refract Surg 2004;30:675–84.

[18] Liu X, Yi C. Analysis of the reasons for not undergoing laser corneal refractive surgery in 611 patients. Cell Biochem Biophys 2011;61:711–4.

[19] Baudouin C, Messmer EM, Aragona P, et al. Revisiting the vicious circle of dry eye disease : a focus on the pathophysiology of meibomian gland dysfunction. Br J Ophthalmol 2016;100:300–6.

[20] Sun CC, Chang CK, Ma DH, et al. Dry eye after LASIK with a femtosecond laser or a mechanical microkeratome. Optom Vis Sci 2013;90:1048–56.

[21] Sharma B, Soni D, Saxena H, et al. Impact of corneal refractive surgery on the precorneal tear film. Indian J Ophthalmol 2020;68:2804–12.

[22] Benítez-Del-Castillo JM, Acosta MC, Wassfi MA, et al. Relation between corneal innervation with confocal microscopy and corneal sensitivity with noncontact esthesiometry in patients with dry eye. Invest Ophthalmol Vis Sci 2007;48:173–81.

[23] Shtein RM. Post-LASIK dry eye. Expert Rev Ophthalmol 2011;6:575–82.

[24] Yang LWY, Mehta JS, Liu YC. Corneal neuromediator profiles following laser refractive surgery. Neural Regen Res 2021;16:2177–83.

[25] Lee HK, Lee KS, Kim HC, et al. Nerve growth factor concentration and implications in photorefractive keratectomy vs laser in situ keratomileusis. Am J Ophthalmol 2005;139:965–71.

[26] Li M, Zhao J, Shen Y, et al. Comparison of dry eye and corneal sensitivity between small incision lenticule extraction and femtosecond LASIK for myopia. PLoS One 2013;8, e77797.

[27] Denoyer A, Landman E, Trinh L, et al. Dry eye disease after refractive surgery : comparative outcomes of small incision lenticule extraction versus LASIK. Ophthalmology 2014.

[28] Wong AHY, Cheung RKY, Kua WN, et al. Dry eyes after SMILE. Asia Pac J Ophthalmol (Phila) 2019;8:397–405.

[29] Liu YC, Yam GH, Lin MT, et al. Comparison of tear proteomic and neuromediator profiles changes between small incision lenticule extraction (SMILE) and femtosecond laser-assisted in-situ keratomileusis (LASIK). J Adv Res 2021;29:67–81.

[30] Kung JS, Sales CS, Manche EE. Corneal sensation and dry eye symptoms after conventional versus inverted side-cut femtosecond LASIK : a prospective randomized study. Ophthalmology 2014;121:2311–6.

[31] Donnenfeld ED, Solomon K, Perry HD, et al. The effect of hinge position on corneal sensation and dry eye after LASIK. Ophthalmology 2003 ; 110 : 1023-9 ; discussion 9-30.

[32] Mian SI, Shtein RM, Nelson A, Musch DC. Effect of hinge position on corneal sensation and dry eye after laser in situ keratomileusis using a femtosecond laser. J Cataract Refract Surg 2007;33:1190–4.

[33] Ma KK, Manche EE. Corneal sensitivity and patient-reported dry eye symptoms in a prospective randomized contralateral-eye trial comparing laser in situ keratomileusis and small incision lenticule extraction. Am J Ophthalmol 2022;241:248–53.

[34] Wang B, Naidu RK, Chu R, et al. Dry eye disease following refractive surgery : a 12-month follow-up of SMILE versus FS-LASIK in high myopia. J Ophthalmol 2015;2015, 132417.

[35] Tsai T, Alwees M, Rost A, et al. Changes of subjective symptoms and tear film biomarkers following femto-LASIK. Int J Mol Sci 2022;23.

[36] Labetoulle M, Benitez-Del-Castillo JM, Barabino S, et al. Artificial tears : biological role of their ingredients in the management of dry eye disease. Int J Mol Sci 2022;23.

[37] Albietz JM, Lenton LM, McLennan SG, Earl ML. A comparison of the effect of refresh plus and bion tears on dry eye symptoms and ocular surface health in myopic LASIK patients. CLAO J 2002;28:96–100.

[38] Hazarbassanov RM, Queiroz-Hazarbassanov NGT, Barros JN, Gomes JAP. Topical osmoprotectant for the management of postrefractive surgery-induced dry eye symptoms : a randomised controlled double-blind trial. J Ophthalmol 2018;2018:4324590.

[39] Wallerstein A, Jackson WB, Chambers J, et al. Management of post-LASIK dry eye : a multicenter randomized comparison of a new multi-ingredient artificial tear to carboxymethylcellulose. Clin Ophthalmol 2018;12:839–48.

[40] Panova IE, Titov AV, Mirsaitova DR. [The effects of a trehalose-based drug used after laser keratorefractive surgeries]. Vestn Oftalmol 2020 ; 136 : 110-6.

[41] Wang T, Di Y, Li Y. Combination therapy with 3 % diquafosol tetrasodium ophthalmic solution and sodium hyaluronate : an effective therapy for patients with dry eye after femtosecond laser-assisted in situ keratomileusis. Front Med (Lausanne) 2023;10:1160499.

[42] Liu Y, Qian Y, Li M, et al. Combination therapy with diquafosol sodium and sodium hyaluronate in eyes with dry eye disease after small incision lenticule extraction. In vivo 2023;37:2829–34.

[43] Toda I, Ide T, Fukumoto T, et al. Combination therapy with diquafosol tetrasodium and sodium hyaluronate in patients with dry eye after laser in situ keratomileusis. Am J Ophthalmol 2014;157:616–22.e1.

[44] Mori Y, Nejima R, Masuda A, et al. Effect of diquafosol tetrasodium eye drop for persistent dry eye after laser in situ keratomileusis. Cornea 2014;33:659–62.

[45] Wu Y, Xu L, Song Y, et al. Management of post-LASIK dry eye with intense pulsed light in combination with 0.1 % sodium hyaluronate and heated eye mask. Ophthalmol Ther 2022;11:161–76.
[46] Astakhov YS, Astakhov SY, Lisochkina AB. Assessment of dry eye signs and symptoms and ocular tolerance of a preservative-free lacrimal substitute (Hylabak(R)) versus a preserved lacrimal substitute (Systane(R)) used for 3 months in patients after LASIK. Clin Ophthalmol 2013;7:2289–97.
[47] Sanchez MA, Arriola-Villalobos P, Torralbo-Jimenez P, et al. The effect of preservative-free HP-Guar on dry eye after phacoemulsification : a flow cytometric study. Eye (Lond) 2010;24:1331–7.
[48] Rosenfeld SI. Evaluation and management of post-LASIK dry eye syndrome. Int Ophthalmol Clin 2010;50:191–9.
[49] Iqbal MS, Latif K. Topical cyclosporine in treatment of post-LASIK symptomatic dry eye. PakJOphthalmol 2020;36:381–5.
[50] Salib GM, McDonald MB, Smolek M. Safety and efficacy of cyclosporine 0.05 % drops versus unpreserved artificial tears in dry-eye patients having laser in situ keratomileusis. J Cataract Refract Surg 2006;32:772–8.
[51] Kanellopoulos AJ. Incidence and management of symptomatic dry eye related to LASIK for myopia, with topical cyclosporine A. Clin Ophthalmol 2019;13:545–52.
[52] Torricelli AA, Santhiago MR, Wilson SE. Topical cyclosporine a treatment in corneal refractive surgery and patients with dry eye. J Refract Surg 2014;30:558–64.
[53] Eskina EN, Maychuk NV, Parshina VA, Kukleva OY. [Medicamental correction of pathomorphological changes of the ocular surface in patients with steroid therapy intolerance after photorefractive keratectomy]. Vestn Oftalmol 2019 ; 135 : 67-77.
[54] Noda-Tsuruya T, Asano-Kato N, Toda I, Tsubota K. Autologous serum eye drops for dry eye after LASIK. J Refract Surg 2006;22:61–6.
[55] Metheetrairut C, Ngowyutagon P, Tunganuntarat A, et al. Comparison of epitheliotrophic factors in platelet-rich plasma versus autologous serum and their treatment efficacy in dry eye disease. Sci Rep 2022;12:8906.
[56] Lozano-Sanroma J, Barros A, Alcalde I, et al. Impact of plasma rich in growth factors (PRGF) eye drops on ocular redness and symptomatology in patients with dry eye disease. Medicina (Kaunas) 2023;59.
[57] Alio JL, Rodriguez AE, Abdelghany AA, Oliveira RF. Autologous platelet-rich plasma eye drops for the treatment of post-LASIK chronic ocular surface syndrome. J Ophthalmol 2017;2017:2457620.
[58] Sanchez-Avila RM, Merayo-Lloves J, Fernandez ML, et al. Plasma rich in growth factors for the treatment of dry eye after LASIK surgery. Ophthalmic Res 2018;60:80–6.
[59] Wang C, Peng Y, Pan S, Li L. Effect of insulin-like growth factor-1 on corneal surface ultrastructure and nerve regeneration of rabbit eyes after laser in situ keratomileusis. Neurosci Lett 2014;558:169–74.
[60] Gong Q, Zhang S, Jiang L, et al. The effect of nerve growth factor on corneal nerve regeneration and dry eye after LASIK. Exp Eye Res 2021;203, 108428.
[61] Joo MJ, Yuhan KR, Hyon JY, et al. The effect of nerve growth factor on corneal sensitivity after laser in situ keratomileusis. Arch Ophthalmol 2004;122:1338–41.
[62] Fakih D, Migeon T, Moreau N, et al. Transient receptor potential channels : important players in ocular pain and dry eye disease. Pharmaceutics 2022;14.
[63] Fakih D, Baudouin C, Reaux-Le Goazigo A, Melik Parsadaniantz S. TRPM8 : a therapeutic target for neuroinflammatory symptoms induced by severe dry eye disease. Int J Mol Sci 2020 ; 21.
[64] Fakih D, Guerrero-Moreno A, Baudouin C, et al. Capsazepine decreases corneal pain syndrome in severe dry eye disease. J Neuroinflammation 2021 ;18:111.
[65] Moreno-Montanes J, Bleau AM, Jimenez AI. Tivanisiran, a novel siRNA for the treatment of dry eye disease. Expert Opin Investig Drugs 2018 ;27:421–6.
[66] Huang B, Mirza MA, Qazi MA, Pepose JS. The effect of punctal occlusion on wavefront aberrations in dry eye patients after laser in situ keratomileusis. Am J Ophthalmol 2004 ;137:52–61.
[67] Nemet A, Mimouni M, Hecht I, et al. Post laser-assisted in-situ keratomileusis dry eye disease and temporary punctal plugs. Indian J Ophthalmol 2020 ;68:2960–3.
[68] Alfawaz AM, Algehedan S, Jastaneiah SS, et al. Efficacy of punctal occlusion in management of dry eyes after laser in situ keratomileusis for myopia. Curr Eye Res 2014 ;39:257–62.
[69] Fouda SM, Mattout HK. Comparison between botulinum toxin A injection and lacrimal punctal plugs for the control of post-LASIK dry eye manifestations : a prospective study. Ophthalmol Ther 2017 ;6:167–74.
[70] Mian SZ, Agranat JS, Jacobs DS. Prosthetic Replacement of the Ocular Surface Ecosystem (PROSE) treatment for complications after LASIK. Eye Contact Lens 2016 ;42:371–3.
[71] Di Pascuale MA, Liu TS, Trattler W, Tseng SC. Lipid tear deficiency in persistent dry eye after laser in situ keratomileusis and treatment results of new eye-warming device. J Cataract Refract Surg 2005 ;31:1741–9.
[72] Schallhorn CS, Schallhorn JM, Hannan S, Schallhorn SC. Effectiveness of an eyelid thermal pulsation procedure to treat recalcitrant dry eye symptoms after laser vision correction. J Refract Surg 2017 ;33:30–6.
[73] Shetty R, Khamar P, Nair AP, et al. Assessing clinical and molecular outcomes of prophylactic thermal pulsation therapy on ocular surface health following refractive surgery. Indian J Ophthalmol 2023 ;71:1508–16.
[74] Zhou X, Shen Y, Shang J, Zhou X. Effects of warm compress on tear film, blink pattern and Meibomian gland function in dry eyes after corneal refractive surgery. BMC Ophthalmol 2021 ;21:330.
[75] Fuentes Páez G, Soler Tomas JR, Burillo S. Intense pulsed light : results in chronic dry eye syndrome after LASIK. Arch Soc Esp Oftalmol (Engl Ed) 2020 ;95:226–30.
[76] Pazo EE, Huang H, Fan Q, et al. Intense pulse light for treating post-LASIK refractory dry eye. Photobiomodul Photomed Laser Surg 2021 ;39:155–63.
[77] Brignole-Baudouin F, Baudouin C, Aragona P, et al. A multicentre, double-masked, randomized, controlled trial assessing the effect of oral supplementation of omega-3 and omega-6 fatty acids on a conjunctival inflammatory marker in dry eye patients. Acta Ophthalmol 2011 ;89:e591–e7.
[78] Giannaccare G, Pellegrini M, Sebastiani S, et al. Efficacy of omega-3 fatty acid supplementation for treatment of dry eye disease : a meta-analysis of randomized clinical trials. Cornea 2019 ;38:565–73.
[79] Asbell PA, Maguire MG, Pistilli M, et al. n-3 Fatty acid supplementation for the treatment of dry eye disease. N Engl J Med 2018 ;378:1681–90.
[80] Esquenazi S, Bazan HE, Bui V, et al. Topical combination of NGF and DHA increases rabbit corneal nerve regeneration after photorefractive keratectomy. Invest Ophthalmol Vis Sci 2005 ;46:3121–7.
[81] He J, Bazan HE. Omega-3 fatty acids in dry eye and corneal nerve regeneration after refractive surgery. Prostaglandins Leukot Essent Fatty Acids 2010 ;82:319–25.
[82] Goyal P, Jain AK, Malhotra C. Oral omega-3 fatty acid supplementation for laser in situ keratomileusis-associated dry eye. Cornea 2017 ;36:169–75.
[83] Galor A, Patel S, Small LR, et al. Pregabalin failed to prevent dry eye symptoms after laser-assisted in situ keratomileusis (LASIK) in a randomized pilot study. J Clin Med 2019 ;8.
[84] Paik DW, Lim DH, Chung TY. Effects of taking pregabalin (Lyrica) on the severity of dry eye, corneal sensitivity and pain after laser epithelial keratomileusis surgery. Br J Ophthalmol 2022 ;106:474–9.
[85] Ghaffariyeh A, Chamacham T. Tricyclic antidepressants : potential therapeutic alternatives for treatment of dry eye symptoms after LASIK. J Refract Surg 2008 ;24:770–1. author reply 1-2.
[86] Labetoulle M, Baudouin C, Calonge M, et al. Role of corneal nerves in ocular surface homeostasis and disease. Acta Ophthalmol 2018.
[87] Galor A, Hamrah P, Haque S, et al. Understanding chronic ocular surface pain : an unmet need for targeted drug therapy. Ocul Surf 2022 ;26:148–56.
[88] Lee JH, Han K, Kim TH, et al. Acupuncture for dry eye syndrome after refractive surgery : a randomized controlled pilot trial. Integr Med Res 2021 ;10, 100456.