Bibliographie

[1] Rapport de l'Agence de la biomédecine 2023.

[2] Gain P, Jullienne R, He Z, et al. Global survey of corneal transplantation and eye banking. JAMA Ophthalmol 2016;134(2):167–73.

[3] Zanetti E, Bruni A, Mucignat G, et al. Bacterial contamination of human organ-cultured corneas. Cornea 2005;24(5):603–7.

[4] Thuret G, Carricajo A, Chiquet C, et al. Sensitivity and rapidity of blood culture bottles in the detection of cornea organ culture media contamination by bacteria and fungi. Br J Ophthalmol 2002;86(12):1422–7.

[5] Thuret G, Carricajo A, Vautrin AC, et al. Efficiency of blood culture bottles for the fungal sterility testing of corneal organ culture media. Br J Ophthalmol 2005;89(5):586–90.

[6] Delbosc B, Hervé P, Carbillet JP, Montard M. Proposal for a medium for short-term preservation of human cornea. Bull Soc Ophtalmol Fr 1983;83(2):217–9.

[7] Piquot X, Delbosc B, Herve P, et al. Preservation of human corneas in organ culture : results of a feasibility clinical protocol. Bull Soc Ophtalmol Fr 1990;90(4):429–32.

[8] Albon J, Tullo AB, Aktar S, Boulton ME. Apoptosis in the endothelium of human corneas for transplantation. Invest Ophthalmol Vis Sci 2000;41(10):2887–93.

[9] Gain P, Thuret G, Chiquet C, et al. Value of two mortality assessment techniques for organ cultured corneal endothelium : trypan blue versus TUNEL technique. Br J Ophthalmol 2002;86(3):306–10.

[10] Thuret G, Chiquet C, Bernal F, et al. Prospective, randomized clinical and endothelial evaluation of 2 storage times for cornea donor tissue in organ culture at 31 degrees C. Arch Ophthalmol Chic Ill 1960 2003;121(4):442–50.

[11] Borderie VM, Baudrimont M, Lopez M, et al. Evaluation of the deswelling period in dextran-containing medium after corneal organ culture. Cornea 1997;16(2):215–23.

[12] Thuret G, Manissolle C, Campos-Guyotat L, et al. Animal compound-free medium and poloxamer for human corneal organ culture and deswelling. Invest Ophthalmol Vis Sci 2005;46(3):816–22.

[13] Zhao M, Thuret G, Piselli S, et al. Use of poloxamers for deswelling of organ-cultured corneas. Invest Ophthalmol Vis Scir 2008;49(2):550–9.

[14] Borderie V, Sabolic V, Touzeau O, et al. Screening human donor corneas during organ culture for the presence of guttae. Br J Ophthalmol 2001;85(3):272–6.

[15] Flury M, He Z, Campolmi N, et al. Fabrication of optical mosaics mimicking human corneal endothelium for the training and assessment of eye bank technicians. Opt Lett 2012;37(1):22–4.

[16] Nejepinska J, Juklova K, Jirsova K. Organ culture, but not hypothermic storage, facilitates the repair of the corneal endothelium following mechanical damage. Acta Ophthalmol (Copenh) 2010;88(4):413–9.

[17] Frueh BE, Böhnke M. Prospective, randomized clinical evaluation of Optisol vs organ culture corneal storage media. Arch Ophthalmol Chic Ill 1960 2000;118(6):757–60.

[18] Delbosc B, Naegelen J, Herve P, et al. Preservation of human corneas in an enriched culture medium at +37 degrees centigrade : histological and biochemical analyses. J Fr Ophtalmol 1987;10(10):547–9. 551-5.

[19] Borderie VM, Laroche L. Microbiologic study of organ-cultured donor corneas. Transplantation 1998;66(1):120–3.

[20] Gain P, Thuret G, Chiquet C, et al. Use of a pair of blood culture bottles for sterility testing of corneal organ culture media. Br J Ophthalmol 2001;85(10):1158–62.

[21] Hermel M, Salla S, Hamsley N, et al. Detection of contamination during organ culture of the human cornea. Graefes Arch Clin Exp Ophthalmol 2010;248(1):117–26.

[22] Utheim TP, Raeder S, Utheim ØA, et al. Sterility control and long-term eye-bank storage of cultured human limbal epithelial cells for transplantation. Br J Ophthalmol 2009;93(7):980–3.

[23] McCarey BE, Edelhauser HF, Van Horn DL. Functional and structural changes in the corneal edothelium during in vitro perfusion. Invest Ophthalmol 1973;12(6):410–7.

[24] Zhao B, Cooper LJ, Brahma A, et al. Development of a three-dimensional organ culture model for corneal wound healing and corneal transplantation. Invest Ophthalmol Vis Sci 2006;47(7):2840–6.

[25] Brunette I, Nelson LR, Bourne WM. A system for long-term corneal perfusion. Invest Ophthalmol Vis Sci 1989;30(8):1813–22.

[26] Thiel MA, Morlet N, Schulz D, et al. A simple corneal perfusion chamber for drug penetration and toxicity studies. Br J Ophthalmol 2001;85(4):450–3.

[27] Tanelian DL, Bisla K. A new in vitro corneal preparation to study epithelial wound healing. Invest Ophthalmol Vis Sci 1992;33(11):3024–8.

[28] Garcin T, Gauthier AS, Crouzet E, et al. Innovative corneal active storage machine for long-term eye banking. Am J Transplant 2019;19(6):1641–51.

[29] Garcin T, Gauthier AS, Crouzet E, et al. Three-month storage of human corneas in an active storage machine. Transplantation 2020;104(6):1159–65.

[30] Regnier M, Auxenfans C, Maucort-Boulch D, et al. Eye bank prepared versus surgeon cut endothelial graft tissue for Descemet membrane endothelial keratoplasty : an observational study. Medicine (Baltimore) 2017;96(19), e6885.

[31] Ragunathan S, Ivarsen A, Nielsen K, Hjortdal J. Comparison of organ cultured precut corneas versus surgeon-cut corneas for Descemet's stripping automated endothelial keratoplasty. Cell Tissue Bank 2014;15(4):573–8.