

BIBLIOGRAPHIE

74.1. Traitement oculoplastique des pathologies de la surface

- [1] Robert PY, Adenis JP. Anatomie des paupières. In: Adenis JP, editor. Chirurgie palpébrale. Paris: Masson; 2007. p. 1–15.
- [2] Adenis JP. Troubles de la statique palpébrale. In: Adenis JP, editor. Chirurgie palpébrale. Paris: Masson; 2007. p. 18–67.
- [3] Robert PY, Adenis JP. Symblépharons. In: Adenis JP, editor. Chirurgie palpébrale. Paris: Masson; 2007. p. 147–9.
- [4] Huang FM, Dong BS, Zhao JF, Hu ZC. A conjunctival pulling-back technique for treatment of serious symblepharon. *Zhonghua Zheng Xing Wai Ke Za Zhi* 2003;19:118–9.
- [5] Orth B, Press UP, Hubner H, Pleyer U. Transplantation of autologous oral mucosa in the treatment of a symblepharon in Wegener's disease. A case report *Klin Monatsbl Augenheilkd* 2001;218:514–7.
- [6] Shree N, Das S, Arya D, et al. Single-staged surgical correction of eyelid sequelae along with lid margin mucous membrane grafting in Stevens-Johnson syndrome and other cicatricial ocular surface diseases. *Cornea* 2023;42:404–11.
- [7] Solomon A, Espana EM, Tseng SC. Amniotic membrane transplantation for reconstruction of the conjunctival fornices. *Ophthalmology* 2003;110:93–100.
- [8] Zhou SY, Chen JQ, Chen LS, et al. Long-term results of amniotic membrane transplantation for conjunctival surface reconstruction. *Zhonghua Yan Ke Za Zhi* 2004;40:745–9.
- [9] Jain S, Rastogi A. Evaluation of the outcome of amniotic membrane transplantation for ocular surface reconstruction in symblepharon. *Eye* 2004;18:1251–7.
- [10] Honavar SG, Bansal AK, Sangwan VS, Rao GN. Amniotic membrane transplantation for ocular surface reconstruction in Stevens-Johnson syndrome. *Ophthalmology* 2000;107:975–9.
- [11] Barabino S, Rolando M. Amniotic membrane transplantation elicits goblet cell repopulation after conjunctival reconstruction in a case of severe ocular cicatricial pemphigoid. *Acta Ophthalmol Scand* 2003;81:68–71.
- [12] Barabino S, Rolando M, Bentivoglio G, et al. Role of amniotic membrane transplantation for conjunctival reconstruction in ocular-cicatricial pemphigoid. *Ophthalmology* 2003;110:474–80.
- [13] Komai S, Inatomi T, Nakamura T, et al. Long-term outcome of cultivated oral mucosal epithelial transplantation for fornix reconstruction in chronic cicatrizing diseases. *Br J Ophthalmol* 2022;106(10):1355–62.
- [14] Zhao D, Yin H, Cheng A, et al. Sealing of the gap between the conjunctiva and tenon capsule to improve symblepharon surgery. *Am J Ophthalmol* 2015;160:438–46.
- [15] Geerling G, Yaici R, Roth M. Treatment of ocular mucous membrane pemphigoid. *Ophthalmology* 2023;120:496–501.

[16] Ollero R, Soldevila A, Fideliz de la Paz M, Martínez-Grau A. Surgical management of ocular cicatricial pemphigoid in a pediatric patient. *Cornea* 2022;41:911–3.

[17] Hokyung Choung H, Reshef E, Tanking T, Freitag S. A conjunctival-sparing surgical technique for lower eyelid cicatricial entropion repair in ocular cicatricial pemphigoid. *Orbit* 2020;39:23–30.

[18] Satchi K, McNab A. Conjunctival cicatrizing disease presenting with lacrimal obstruction. *Orbit* 2016;35:321–3.

[19] Juniat V, Joshi S, Hersch D, Selva D, Joshi N. Eye Canthoplasty repair for canthal rounding. *Eye* 2022;36:564–7.

74.2. Complications cornéennes de la chirurgie oculoplastique

[20] Bagheri A, Najmi H, Salim RE, Yazdani S. Tear condition following unilateral ptosis surgery. *Orbit* 2015;34(2):66–71.

[21] George JL, Tercero ME, Angioi-Duprez K, Maalouf T. Risk of dry eye after mullerectomy via the posterior conjunctival approach for thyroid-related upper eyelid retraction. *Orbit* 2002;21(1):19–25.

[22] Gonnermann J, Klein JP, Klamann MKJ, et al. Dry eye symptoms in patients after eyelid reconstruction with full-thickness eyelid defects : using the Tomey TG-1000 thermographer. *Ophthalmic Res* 2012;48(4):192–8.

[23] Kim HH, De Paiva CS, Yen MT. Effects of upper eyelid blepharoplasty on ocular surface sensation and tear production. *Can J Ophthalmol* 2007;42(5):739–42.

[24] <https://www.sofcprefr.fr/src/fiches-informations/fr/chirurgie-esthetique-des-paupieres-ou-blepharoplastie-esthetique.pdf>.

[25] Parikh M, Kwon YH. Vision loss after inadvertent corneal perforation during lid anesthesia. *Ophthalmic Plast Reconstr Surg* 2011;27(5):e141–2.

[26] Shoji MK, Tran AQ, Nikpoor N, Lee WW. Corneal laceration associated with upper eyelid blepharoplasty. *Ophthalmic Plast Reconstr Surg* 2020;36(1):e21–3.

[27] Maalouf T, George JL. Serious corneal complication after ptosis surgery : six case reports. *J Fr Ophthalmol* 2007;30(9):893–8.

[28] Chen PJ, Lee YK, Lai CC. Significant changes of corneal astigmatism after levator muscle surgery for acquired blepharoptosis. *Ophthalmic. Plast Reconstr Surg* 2024. Online ahead of print.

[29] Francisco ZM, Carlos R de L, Santiago OP, et al. Refractive changes associated with oculoplastic surgeries : A scoping review. *Eur J Ophthalmol* 2024. 11206721241247422.

[30] Zinkernagel MS, Ebnetter A, Ammann-Rauch D. Effect of upper eyelid surgery on corneal topography. *Arch Ophthalmol* 2007;125(12):1610–2.

[31] Saleh GM, Mavrikakis I, de Sousa JL, et al. Corneal astigmatism with upper eyelid gold weight implantation using the combined high pretarsal and levator fixation technique. *Ophthalmic Plast Reconstr Surg* 2007;23(5):381–3.